

## Tilburg University

### Automata, matching and foraging behavior of bees

Thuijsman, F.; Peleg, B.; Amitai, M.; Shmida, A.

*Publication date:*  
1993

[Link to publication in Tilburg University Research Portal](#)

*Citation for published version (APA):*

Thuijsman, F., Peleg, B., Amitai, M., & Shmida, A. (1993). *Automata, matching and foraging behavior of bees*. (CentER Discussion Paper; Vol. 1993-78). CentER.

#### General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

#### Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

CBM  
8414 R  
993-78  
8414  
1993  
78  
ntER  
for  
nic Research

# Discussion paper



Center  
for  
Economic Research

No. 9378

*Ru*  
*Environment*  
*mathematics*

**AUTOMATA, MATCHING AND  
FORAGING BEHAVIOR OF BEES**

by F. Thuijsman, B. Peleg,  
M. Amitai and A. Shmida

December 1993

ISSN 0924-7815



K.U.B.

BIBLIOTHEEK  
TILBURG



# Automata, Matching and Foraging Behavior of Bees<sup>1</sup>

F. Thuijsman<sup>2</sup>, B. Peleg<sup>3</sup>, M. Amitai<sup>3</sup> & A. Shmida<sup>4</sup>

October 1993

## Abstract

We discuss two types of simple foraging strategies for bees. Each of these explicit strategies explains that in a multi-bee community the bees will distribute themselves over the nectar sources according to the Ideal Free Distribution. At the same time these strategies explain that in single-bee experimental settings a bee will match, by its number of visits, the nectar supply from the available sources (the Matching Law). Moreover, both strategies explain that in certain situations the bees may behave as if they are risk averse. Our results indicate that a competitive market in a multi-bee community permits individuals to be boundedly rational and still forage optimally.

---

<sup>1</sup>This is a revision of discussion paper 30, Center for Rationality, Hebrew University of Jerusalem, Israel

<sup>2</sup>Department of Mathematics, University of Limburg, Maastricht, The Netherlands

<sup>3</sup>Department of Mathematics and Center for Rationality, The Hebrew University of Jerusalem, Israel

<sup>4</sup>Department of Evolution and Ecology and Center for Rationality, The Hebrew University of Jerusalem, Israel

# 1 Introduction

In animal behavior studies many experiments of the following type have been done (Commons et al. 1982, Maynard-Smith 1982, Staddon 1983, Houston 1983, Davison & McCarthy 1988, Menzel & Gregger 1992). An animal subject can repeatedly choose from two different food sources that give precisely the same quantity of the same food, however with different frequencies. Suppose that, at each point in time, the yellow source  $Y$  gives one unit of food with probability  $p$ , and the blue source  $B$  gives one unit of food with probability  $q$  (and 0 units otherwise). These experiments reveal that the animal subject behaves as if, asymptotically, it matches the reward probabilities by the frequencies of visiting the alternatives, i.e. the number of visits to  $Y$  relates to the number of visits to  $B$  as  $p$  to  $q$  (cf. Staddon 1983, Houston & Sumida 1987, Krebs & Kacelnik 1991, Menzel & Gregger 1992). This kind of behavior has been called “the Matching Law” (see for example Houston 1983, Krebs & Kacelnik 1991).

These single-animal experiments are closely related to so-called two-armed bandit problems in statistical decision theory, where a player wishes to maximize his profits from playing a two-armed gambling machine of which one arm gives a greater probability of winning. The rational player, not knowing the winning probabilities of the arms, should try to find the more profitable arm and play it exclusively (DeGroot 1970, Rothschild 1974). A behavior of matching the success probabilities seems to be irrational (Houston et al. 1982, Staddon 1983, Real 1992). One of our goals is to explain this “irrational” behavior by presenting simple foraging strategies that are optimal and rational in a multi-animal natural environment.

We show that in a multi-animal setting these strategies lead to what is known as the Ideal Free Distribution (IFD, Fretwell 1972, Milinski & Parker 1991). Peleg & Shmida (1992) have proved the existence of the IFD in a general framework of an environment with many flower types and

bee types. To explain briefly the IFD, consider  $n$  (identical) animals and two food sources  $Y$  and  $B$  where food is being supplied with rates  $p$  and  $q$  respectively. Then after some time a fraction  $p/(p+q)$  of the  $n$  animals will be feeding at the  $Y$  source while a fraction of  $q/(p+q)$  will be at  $B$ . Thus, the number of animals at  $Y$  relates to the number of animals at  $B$  as  $p$  to  $q$ . In a natural multi-animal situation, establishing the IFD seems to be the rational thing to do, since at the IFD the average intake of food is the same at all food sources. The population is at equilibrium; no animal can improve its payoff by feeding at another food source. When non-cooperatively trying to establish the IFD is the "natural" thing to do for the animal, then in an artificial single-animal setting one may observe irrational behavior.

When applying our strategies to a situation where the animal can choose between two probabilistic food sources in which the food supply is normally distributed with the same mean but with different variances, our result shows that the animal will act as if it is risk averse or, as if it is risk prone (sensu Real & Caraco 1986). An animal will act as if it is risk averse if its (individual) critical level is smaller than the mean, and risk prone otherwise. Here the critical level is a threshold by which the animal judges each quantity of food it obtains as being satisfactory or not. The animal will prefer the food source with the smallest probability of getting something below critical level.

For simplicity we shall talk about bees instead of animals in this paper. Most of the wild bees, especially in the Mediterranean area, are solitary bees which live about four-five weeks (Shmida et al. 1993). Each female has its own nest and raises its brood alone. There is no exchange of foraging information among bees. Each bee is assumed to be a maximizer of Darwinian fitness (Hammerstein & Selten 1993), i.e. it maximizes its average nectar intake per time unit. Nectar will be considered as the only reward for

the bee in the model. Nectar is produced by flowers for the sole purpose of attracting pollinators. Bees have no innate preference to particular flowers (Heinrich 1979, Menzel 1985, 1990, Menzel & Shmida 1992) and their foraging patterns appear to be influenced by experience. Experiments reveal that bees use mainly their short-term memory for decision-making in local flower patches (Menzel & Gregger 1992) and take into account only the last one or two flowers in the process of deciding on leaving or staying in a patch of a given flower type (Real et al. 1990, Cresswell 1990, Real 1991, 1992, Kadmon & Shmida 1992, Kadmon et al. 1992). The above experimental result initiated our approach to assume that the bees have bounded recall. The strategies we provide are described by finite automata (Ben-Porath & Peleg 1987) by which the bees respond only to their own payoffs and remember only payoffs of the last few visits. In this paper we neither include travel times nor handling times for the flower types. These parameters are considered in some other models (Lavery 1980, Harder 1987, Friedman & Shmida 1992, Peleg & Shmida 1992).

The two explicit strategies examined in this paper shall be called *the  $\epsilon$ -sampling strategy* and *the failures strategy*. Briefly one could say that the  $\epsilon$ -sampling strategy is to visit one alternative repeatedly, but every now and then sampling on the other alternative and to switch if the other alternative is better than the one previously selected. The failures strategy describes an innate behavior leading to matching according to some simple finite automaton (e.g. leave  $Y$  after  $y$  empty flowers, leave  $B$  after  $b$  empty flowers). The two behavioral strategies represent two alternative approaches to model choice rules of foraging animals: resource dependent moves vs. resource independent moves. These strategies will be discussed in sections 2 and 3 respectively, each with respect to multi-animal and single-animal settings and in relation to risk. Section 4 concludes with some discussion on related literature.

To simplify notations we restrict our attention to the situation where there are only two food sources:  $Y$  (yellow) and  $B$  (blue). However, our results can be extended to situations with any finite number of food sources.



## 2 The $\epsilon$ -Sampling Strategy

The  $\epsilon$ -sampling strategy is briefly described as follows. A foraging bee which has to choose between two colored resources will use the following rule: initially choose one of the colors at random, then at each point in time stay at the current color with probability  $1 - \epsilon$  and sample the other color with probability  $\epsilon$ ; when sampling, if you find a payoff above your critical level, then switch to this new color, otherwise return to the previous color immediately; at this next color (new or old) again at each point in time stay with probability  $1 - \epsilon$ , sample elsewhere with probability  $\epsilon$ . This  $\epsilon$ -sampling strategy is close to Heinrich's (1979) idea of "Major-Minor behavior" of bees, which motivated our research.

**Definition 2.1** Let  $\alpha, \epsilon \in (0, 1)$ , let  $a(t) \in \{Y, B\}$  represent the action selected and let  $r(t) \in \mathbb{R}$  be the payoff at time  $t \in \{1, 2, 3, \dots\}$ .

Define  $cl(1) = 0$  and

$$cl(t + 1) = \alpha cl(t) + (1 - \alpha)r(t) \quad (1)$$

for  $t \geq 1$ . Then  $cl(t)$  is called the critical level at time  $t$ .

Let  $Y_\epsilon$  denote the mixed action: choose  $Y$  with probability  $1 - \epsilon$  and  $B$  otherwise and let  $B_\epsilon$  be defined similarly.

The  $\epsilon$ -sampling strategy is defined by playing:

at  $t = 1$  use  $Y_{0.5}$ ,

at  $t = 2$  use  $a(1)_\epsilon$ ,

at  $t > 2$  use  $a(t - 1)_\epsilon$  in case  $a(t - 1) \neq a(t - 2)$  and  $r(t - 1) > cl(t - 1)$ , use  $a(t - 2)_\epsilon$  otherwise.

Note that equation (1) is the linear operator function originally used by Bush & Mosteller (1955). In this definition the parameters  $\alpha$  and  $\epsilon$  are the individual bee's factors. The first one is related to memory and the second one to searching elsewhere. Those familiar with game theory can think of

$\epsilon$  as a bee's *trembling hand* (Selten 1975) trying to play a pure action. We wish to emphasize that different individuals may have different parameters, but our results remain the same.

Let  $\mathbf{N}$  denote the set of natural numbers  $\{1, 2, 3, \dots\}$ . Then, consider what happens if we have a population of  $n \in \mathbf{N}$  bees that are foraging on two patches of flowers, a yellow patch  $Y$  and a blue patch  $B$ . Per unit of time the yellow patch has a total nectar supply of  $y \in \mathbf{R}_+$ , while there is a total quantity of  $b \in \mathbf{R}_+$  at patch  $B$ . We make the following assumptions:

- a. If at some stage  $t \in \mathbf{N}$  there are  $n_Y$  bees at  $Y$ , then we assume that at this stage each of these bees is receiving  $r(t) = y/n_Y$  units of nectar. This means that the total quantity of nectar at  $Y$  is being equally distributed over the visiting bees at  $Y$  at each stage. Likewise for  $B$ .
- b. There is no accumulation of nectar at a patch. All nectar is taken by the bees at each stage and if at some stage there are no bees at a patch then at the next stage the total quantity is still the same.
- c. The bees sampling factors  $\epsilon$  are sufficiently close to 0 to have a negligible probability of two or more bees moving at the same time. Thus the distribution of bees over flower types changes by one bee moving either from  $Y$  to  $B$  or from  $B$  to  $Y$ . (Here "moving from  $Y$  to  $B$ " means: previously the bee was at  $Y$ , now it has gone to  $B$  for a sample and since the payoff received at  $B$  is larger than the critical level it decides to stay at  $B$ .)
- d. We assume that, when going out sampling, the bee has been in  $Y$  sufficiently long to have its critical level close to  $y/n_Y$ .

**Theorem 2.2** *Under the above assumptions the population will stabilize in the Ideal Free Distribution.*

**Proof:** Firstly, observe that, due to its  $\varepsilon$ , each bee will go out sampling the other color infinitely often. Hence, if the process stabilizes in some distribution of bees over  $Y$  and  $B$ , then it must be such that no single bee can strictly improve its payoff by moving to the other color. Secondly, observe that whenever a bee moves from  $Y$  to  $B$  we must have that  $y/n_Y < b/(n_B + 1)$ , where  $n_Y$  and  $n_B$  are the numbers of bees before the move at  $Y$  and  $B$  respectively.

We now define a *potential function* (Monderer & Shapley 1988) on the distributions of bees as

$$P(n_Y, n_B) = y \sum_{m=1}^{n_Y} 1/m + b \sum_{m=1}^{n_B} 1/m, \quad (2)$$

where  $\sum_{m=1}^0 1/m$  is understood to be equal to 0. If a bee moves from  $Y$  to  $B$ , then the distribution changes from  $(n_Y, n_B)$  to  $(n_Y - 1, n_B + 1)$ . At the same time the potential changes from  $P(n_Y, n_B)$  to  $P(n_Y - 1, n_B + 1)$ . Now  $P(n_Y - 1, n_B + 1) - P(n_Y, n_B) = b/(n_B + 1) - y/n_Y > 0$  because the bee decided to stay at  $B$ . Hence, with each bee movement the potential strictly increases. Since there are only finitely many distributions of the  $n$  bees and since the bees will keep moving as long as possible, the potential function will eventually reach its maximum. At this maximum we have that  $y/n_Y \geq b/(n_B + 1)$  and  $b/n_B \geq y/(n_Y + 1)$ ; hence

$$y/n_Y \approx b/n_B \quad (3)$$

and the population has reached the IFD.  $\square$

We remark that the stable situation that arises from all bees doing  $\varepsilon$ -sampling strategies, needs not be Pareto-optimal. Take for example 4 bees, 8 units of nectar at  $Y$  and 1 unit of nectar at  $B$ . For this situation the IFD will be that all 4 bees will go to  $Y$  and the 1 unit at  $B$  is not being consumed. Thus each bee is getting a payoff 2 all the time. The bees could all improve their payoffs by visiting  $B$  in turns, giving each bee the



average payoff  $9/4 > 2$ . For a continuum of bees the distribution obtained from the  $\epsilon$ -sampling strategies will indeed be Pareto-optimal, so that in a natural environment with many bees and flowers the IFD solution obtained is practically Pareto-optimal.

It should be noted that, to derive the IFD result of Theorem 2.2 one can also take the alternative potential function  $P'$  defined by

$$P'(n_Y, n_B) = \min\{y/n_Y, b/n_B\} \quad (4)$$

for  $n_Y \neq 0$  and  $n_B \neq 0$ , while  $P'(0, n_B) = b/n_B$  and  $P'(n_Y, 0) = y/n_Y$ . Using this alternative potential function one can even show that a population of  $\epsilon$ -sampling bees with different sizes will stabilize in the IFD. Here we assume that a bee of size  $i$  consumes  $i$  times as much nectar as a bee of size 1. Correspondingly, one should think of  $n_Y$  and  $n_B$  as the total size of all bees present at  $Y$  and  $B$  respectively. If we have a finite number of bees with sizes in  $[1, k]$ , where  $k$  is the maximum size present, then the population has the IFD if and only if  $y/n_Y \geq b/(n_B + i)$  and  $b/n_B \geq y/(n_Y + j)$ , where  $i$  ( $j$ ) is the smallest size present at  $Y$  (resp.  $B$ ). Notice that, if  $n_Y \neq 0$  and  $y/n_Y < b/(n_B + i)$ , then  $y/(n_Y + j) < y/n_Y < b/(n_B + i) < b/n_B$  implying that no bee will move from  $B$  to  $Y$ . On the other hand, a bee of size  $i$  sampling from  $Y$  at  $B$  would decide to stay at  $B$ . There is at least one bee of size  $i$  at  $Y$  and hence, by the  $\epsilon$ -sampling strategy at least one bee will move from  $Y$  to  $B$ . Suppose that it has size  $x \geq i$ . Then the distribution changes from  $(n_Y, n_B)$  to  $(n_Y - x, n_B + x)$  and  $P'(n_Y - x, n_B + x) = \min\{y/(n_Y - x), b/(n_B + x)\} > y/n_Y = \min\{y/n_Y, b/n_B\} = P'(y/n_Y, b/n_B)$ . Hence, with each moving bee  $P'$  is strictly increasing and the process will reach a (local) maximum after finitely many moves. At this (local) maximum the population is in IFD. Remark that, if all bees have the same size, then every local maximum of this alternative potential function  $P'$ , as of the original potential function  $P$ , is necessarily a global maximum.




Figure 1: Markov chain for the  $\varepsilon$ -sampling strategy

Let us now see what happens if we take a single bee for an experiment of the following type (Figure 1). We have two artificial *Bernoulli flowers*  $Y$  (yellow) and  $B$  (blue). Each time the bee visits the yellow flower it will receive 1 unit of nectar with probability  $p$  and 0 units otherwise. Equivalently, in a natural situation one can think of a patch of yellow flowers, where a  $p$ -fraction of the flowers is full, while a  $(1 - p)$ -fraction is empty. For the blue flower we have probability  $q$  for a full flower. Let, as before,  $\varepsilon$  be the bee's sampling factor. We assume that when sampling from  $Y$  to  $B$  the bee's critical level will approximately be  $p$  and when sampling from  $B$  to  $Y$  it will approximately be  $q$ . If the bee applies the  $\varepsilon$ -sampling strategy, then its behavior corresponds with the Markov chain depicted in Figure 1. In this figure  $Y_1$  and  $B_1$  correspond to the sampling stages at the respective colors; one gets there by sampling probability  $\varepsilon$  and one decides to remain (or else to return) with probability  $p$  and  $q$  respectively. Consequently,  $Y_2$  and  $B_2$  represent situations where the bee has decided to stay in the particular color.

Computing the stationary distribution of this Markov chain, one finds that the number of visits to  $Y$  relates to the number of visits to  $B$  as  $p + q\varepsilon$

to  $q + p\varepsilon$ , that is approximately as  $p$  to  $q$  if  $\varepsilon$  is small. Thus we have the following theorem.

**Theorem 2.3** *A bee applying the  $\varepsilon$ -sampling strategy in a single-bee experiment will exhibit matching the payoff probabilities by the frequency of its visits (the Matching Law).*

## Attitude Towards Risk

Consider a single bee foraging on two patches  $Y$  and  $B$ , in which all flowers have a normally distributed nectar supply with a common mean  $\mu$ , but where the distributions differ in variance; for the yellow flowers the variance is  $y$  while for the blue ones it is  $b$ .

**Theorem 2.4** *Suppose that the bee is using the  $\varepsilon$ -sampling strategy and has a fixed critical level  $cl$  to decide whether any flower is full or empty. Without loss of generality we assume  $y < b$ .*

*If  $cl < \mu$ , then the bee will act as if it is risk averse, i.e. it will spend more time on  $Y$  than on  $B$ .*

*If  $cl > \mu$ , then the bee will act as if it is risk prone, i.e. it will spend more time on  $B$  than on  $Y$ .*

**Proof:** If  $cl < \mu$ , then the probability of getting a reward above the critical level is larger for  $Y$  than it is for  $B$  (cf. Figure 2). So, when sampling, the bee will decide to move to  $Y$  more often than it will decide to move to  $B$ . Hence the result. The second part is similar.  $\square$

This result can be extended straightforwardly to continuous nectar supply distributions  $F_Y, F_B$  on an interval  $[\alpha, \beta]$  which have a common mean  $\mu = (\alpha + \beta)/2$  and continuous derivatives  $f_Y, f_B$  for which

- a.  $f_i(\mu + x) = f_i(\mu - x)$  for all  $x \in [0, \beta - \mu]$  and  $i = Y, B$  (which means that  $f_i$  is symmetric with respect to the mean  $\mu$ ), and




Figure 2: Two patches of flowers with normal nectar distributions

b. there is  $x^* \in [0, \beta - \mu]$  with

$$f_Y(x) > f_B(x) \text{ for all } x \in (\mu - x^*, \mu + x^*),$$

$$f_Y(x) < f_B(x) \text{ for all } x \in [\alpha, \mu - x^*) \cup (\mu + x^*, \beta].$$

It should be noted that the result depends on how  $F_Y(cl)$  and  $F_B(cl)$  are compared. If  $F_Y(cl) < F_B(cl)$ , then an  $\epsilon$ -sampling bee with critical level  $cl$  will favour the yellow patch, since  $F_i(cl)$  is the probability of having an empty flower at  $i = Y, B$ . Thus, if  $F_Y, F_B$  are nectar supply distribution functions on  $(0, \infty)$  with  $F_Y(r) < F_B(r)$  for all  $r \in (0, \infty)$  (i.e.  $F_Y$  strictly dominates  $F_B$  according to the “first order stochastic dominance criterion” (Fishburn 1976)), then *every* bee that uses an  $\epsilon$ -sampling strategy and judges flowers full or empty according to some critical level, will favour the yellow patch. This means that under these conditions, every bee would visit more yellow flowers than blue flowers or, in other words, spend more than half of its foraging time at the yellow patch.

One can also take nectar distributions as in Figure 3 (or, e.g. negative exponential distributions), in which most flowers are empty or have very small nectar quantities. Such leptokertic distributions are very common in nature (Shreiber 1993, Boker 1993). In case the bee forages on two flower




Figure 3: Two patches of flowers with decreasing nectar distributions

patches  $Y$  and  $B$  with distributions as in Figure 3, then the bee will compare the two patches. If we assume that  $f_Y$  and  $f_B$  are continuous and cross each other only once (e.g.  $f_Y$  starts above  $f_B$ ), then such a comparison also falls in the first order stochastic dominance. If a bee forages on such flower patches and uses a critical level for switching between  $Y$  and  $B$ , then it will stay more frequently in the blue patch with the higher variance than in the yellow patch with the lower variance.

We would like to emphasize that the results presented do not depend on  $\epsilon$ . All that matters is that  $\epsilon \in (0, 1)$  is sufficiently small to have a bee staying in the same patch long enough to get a good estimate of the payoff in this patch. One can even allow  $\epsilon$  to be payoff dependent, getting smaller with high current payoffs and getting larger with low current payoffs. The only thing needed is that no matter at what color (patch) the bee is foraging, with probability 1 it will eventually sample at another color (e.g. this condition is satisfied if  $\epsilon$  is bounded away from 0). In the multi-bee model the IFD result will still be valid if all individual bees have (sufficiently small) different  $\epsilon$ 's.



### 3 The Failures Strategy

In the previous section the bee's "decision" of going out to sample the other color was independent of the payoffs received in the current color; it was determined solely by the innate sampling factor  $\epsilon$ . In this section we present a foraging strategy which will make the bee move to another color after engaging a certain number of consecutive empty flowers. Recall that empty is to be interpreted as "below critical level". The bee is assumed to behave as a finite automaton (Neyman 1985, Ben- Porath & Peleg 1987, Kalai 1988).

This automaton strategy corresponds to the well known "area- restricted search" (Real et al. 1990) of animal behavior, which was termed "near-far" by Motro & Shmida (1992). The near-far strategy is  $A(1, 1)$  (see below), a special case of our automaton strategy where the bee uses only the last flower visited to evaluate the current patch. It means: stay in the patch as long as you find food and leave otherwise. Similar strategies have been reviewed by Houston et al. (1982). Experimental studies reveal this near-far behavior (see in Motro & Shmida 1992).

Let us return to the experiment with two artificial flowers  $Y$  (yellow) and  $B$  (blue) that give 1 unit of nectar with probability  $p$  and  $q$  respectively (and 0 units otherwise). As already mentioned above, observations indicate that the frequencies of visits by the bee will match these probabilities. The event of receiving 1 shall be called a succes, receiving 0 is a failure.

**Definition 3.1** *Let  $y, b \in \mathbb{N}$ . The finite automaton  $A(y, b)$  is given by:*

- a. leave  $Y$  after  $y$  consecutive failures and move to  $B$ ,*
- b. leave  $B$  after  $b$  consecutive failures and move to  $Y$ .*

As an example we have depicted automaton  $A(3, 2)$  in Figure 4-a, where  $Y_k$  indicates the automaton state of being at  $Y$  while the last  $k$  consecutive




Figure 4: (a) Automaton  $A(3,2)$  and (b) related Markov chain

yellow visits were failures ( $k = 0, 1, 2$ ). Obviously  $B_0$  and  $B_1$  are to be interpreted similarly. The automaton corresponds with a Markov chain on the states  $Y_0, Y_1, Y_2, B_0, B_1$ , which is depicted in Figure 4-b. In this figure and elsewhere in this paper we write  $\bar{p}$  to denote  $1 - p$  and  $\bar{q}$  for  $1 - q$ . This Markov chain in turn corresponds to the transition matrix  $T$  given by:

$$T = \begin{matrix} & \begin{matrix} Y_0 & Y_1 & Y_2 & B_0 & B_1 \end{matrix} \\ \begin{matrix} Y_0 \\ Y_1 \\ Y_2 \\ B_0 \\ B_1 \end{matrix} & \begin{pmatrix} p & \bar{p} & 0 & 0 & 0 \\ p & 0 & \bar{p} & 0 & 0 \\ p & 0 & 0 & \bar{p} & 0 \\ 0 & 0 & 0 & q & \bar{q} \\ \bar{q} & 0 & 0 & q & 0 \end{pmatrix} \end{matrix}$$

If the bee uses  $A(3,2)$ , then we can compute its frequencies of visits to  $Y$  and  $B$  by finding the stationary distribution  $\pi = (y_0, y_1, y_2, b_0, b_1)$  of  $T$ . The frequency of visits to  $Y$  is  $y_0 + y_1 + y_2$  and that to  $B$  is  $b_0 + b_1$ . The vector  $\pi$  is nonnegative and adding its components gives 1. Furthermore  $\pi$  has the property that  $\pi T = \pi$ . Using  $A(3,2)$  the bee would confirm the matching

law if and only if

$$\begin{cases} y_0 + y_1 + y_2 = p/(p+q) \\ b_0 + b_1 = q/(p+q) \end{cases} \quad (5)$$

The equation  $\pi T = \pi$  yields

$$y_1 = \bar{p}y_0, y_2 = \bar{p}^2y_0, b_1 = \bar{q}b_0 \text{ and } b_0 = (\bar{p}^3/\bar{q}^2)y_0.$$

Hence we find the equations

$$y_0 + y_1 + y_2 = y_0(1 + \bar{p} + \bar{p}^2) = y_0 \frac{1 - \bar{p}^3}{\bar{p}},$$

$$b_0 + b_1 = b_0(1 + \bar{q}) = b_0 \frac{1 - \bar{q}^2}{\bar{q}} = y_0 \frac{\bar{p}^3(1 - \bar{q}^2)}{\bar{q}^2\bar{q}}.$$

So, if we let  $f(2, 3)$  denote the ratio of the frequencies of visits to  $Y$  and  $B$ , then

$$f(2, 3) = \frac{\text{frequency of visits to } Y}{\text{frequency of visits to } B} = \frac{q\bar{q}^2(1 - \bar{p}^3)}{p\bar{p}^3(1 - \bar{q}^2)} \quad (6)$$

and the automaton  $A(2, 3)$  is matching if and only if  $f(2, 3) = p/q$ , which is equivalent to:

$$\frac{p^2\bar{p}^3}{1 - \bar{p}^3} = \frac{q^2\bar{q}^2}{1 - \bar{q}^2}. \quad (7)$$

Similarly one can show the following theorem.

**Theorem 3.2** *The finite automaton  $A(y, b)$  matches the payoff probabilities  $p$  and  $q$  respectively if and only if*

$$\frac{p^2\bar{p}^y}{1 - \bar{p}^y} = \frac{q^2\bar{q}^b}{1 - \bar{q}^b}. \quad (8)$$

We would like to remark that there are infinitely many  $(y, b) \in \mathbf{R}_+^2$  for which (8) holds, because the function  $y \mapsto \frac{p^2\bar{p}^y}{1 - \bar{p}^y}$  is strictly decreasing on  $(0, \infty)$  from  $\infty$  to 0. In fact we have the following result.

**Theorem 3.3** *Let  $\delta \in (0, 0.5)$  and let  $M \in \mathbf{N}$  be such that  $(1 - \delta)^{M-1} \leq 4\delta(1 - \delta^M)$ . Then, for all  $\delta \leq p \leq q \leq 1 - \delta$  there exist  $y, b \in [1, M]$  with*

$$\frac{p^2\bar{p}^y}{1 - \bar{p}^y} = \frac{q^2\bar{q}^b}{1 - \bar{q}^b}.$$



**Proof:** If  $p\bar{p} = q\bar{q}$ , then  $y = b = 1$  gives the result. Otherwise, assume without loss of generality that  $p\bar{p} < q\bar{q}$ . Then

$$\frac{q^2\bar{q}^M}{1 - \bar{q}^M} = \frac{q\bar{q}q^{M-1}}{(1 - \bar{q}^M)/(1 - \bar{q})} \leq \frac{0.25(1 - \delta)^{M-1}}{(1 - \delta^M)/(1 - \delta)} \leq \delta(1 - \delta) \leq p\bar{p}. \quad (9)$$

By continuity there exists  $b \in [1, M]$  with  $\frac{q^2\bar{q}^b}{1 - \bar{q}^b} = p\bar{p} = \frac{p^2\bar{p}}{1 - \bar{p}}$ .  $\square$

For any  $p, q$ -situation of yellow and blue flowers with  $p, q \in [\delta, 1 - \delta]$  this theorem provides the existence of real numbers  $y, b \in [1, M]$  for which  $\frac{p^2\bar{p}^y}{1 - \bar{p}^y} = \frac{q^2\bar{q}^b}{1 - \bar{q}^b}$ . However, to have an automaton for the bee, we need to have numbers in  $\{1, 2, \dots, M\}$ . We therefore propose to see  $(y, b)$  as a convex combination of pure pairs  $(y^*, b^*) \in \{1, 2, \dots, M\}^2$  and to have the bee perform an initial randomization over the pure automata  $A(y^*, b^*)$  according to the weights of the convex combination. Then, in expectation, the bee will leave  $Y$  after  $y$  consecutive failures and it will leave  $B$  after  $b$  consecutive failures. In view of the bee's bounded recall, it is worthwhile to notice that with  $M = 2$  one can handle  $\delta$  as small as 0.22, while with  $M = 3$  one can handle  $\delta$  as small as 0.18. Thus, for all  $p, q \in [0.22, 0.78]$  one can obtain matching with automata  $A(y, b)$  with  $y, b \in \{1, 2\}$ .

For a bee doing matching with an automaton of above type, the number of consecutive empty flowers it allows itself to encounter in yellow can be different from that in blue. One can think of the bee as having an internal mechanism by which, after some rough estimation of  $p$  and  $q$ , the natural numbers  $y^*$  and  $b^*$  are selected. It is not necessarily true that the bee can use an automaton of type  $A(x, x)$  since there does not always exist a real number  $x \geq 1$  with  $\frac{p^2\bar{p}^x}{1 - \bar{p}^x} = \frac{q^2\bar{q}^x}{1 - \bar{q}^x}$ . If for example one takes  $p = 0.6$  and  $q = 0.8$ , then such  $x$  does not exist. However, we have the following result.

**Theorem 3.4** *If  $p \leq q \leq 1 - p$ , then there exists  $x \geq 1$  such that*

$$\frac{p^2\bar{p}^x}{1 - \bar{p}^x} = \frac{q^2\bar{q}^x}{1 - \bar{q}^x}. \quad (10)$$

**Proof:** If  $p = q$  or if  $\bar{p} = q$ , then one can take  $x = 1$ . Now let  $p < q < \bar{p}$ . Using an automaton  $A(x, x)$  the ratio of the frequencies of visits to  $Y$  and  $B$  is (cf. example  $A(3, 2)$  above)

$$f(x, x) = \frac{q\bar{q}^x(1 - \bar{p}^x)}{p\bar{p}^x(1 - \bar{q}^x)}. \quad (11)$$

Then  $\lim_{x \rightarrow \infty} f(x, x) = 0$  and  $f(1, 1) = \bar{q}/\bar{p} > p/q$ . Hence, by continuity there exists  $x \geq 1$  with  $f(x, x) = p/q$ , or equivalently  $\frac{p^2\bar{p}^x}{1 - \bar{p}^x} = \frac{q^2\bar{q}^x}{1 - \bar{q}^x}$ .  $\square$

Obviously a similar result can be obtained for  $q \leq p \leq \bar{q}$ . Remark that the condition of this theorem is easily satisfied if  $p$  and  $q$  are small. Such conditions are frequently observed in natural situations (Shreiber 1993).

Suppose we have finitely many bees foraging on two patches  $Y$  and  $B$  consisting of Bernoulli flowers. At  $Y$  each flower gives 1 unit of nectar with probability  $p$ . At  $B$  each flower is full with probability  $q$ . We assume that the nectar supply probabilities are independent of the numbers of visiting bees. If in this situation each individual bee determines, by its internal mechanism, an automaton to forage on  $Y$  and  $B$ , then each of these individuals will approximately spend  $\frac{p}{p+q}$  of its foraging time on  $Y$  and  $\frac{q}{p+q}$  on  $B$ . Even stronger, we can say that for each individual bee the probability of being at  $Y$  at time  $t$  converges to  $\frac{p}{p+q}$ . Thus, by the strong law of large numbers the fraction of the numbers of bees we encounter at  $Y$  at time  $t$  converges to  $\frac{p}{p+q}$  as  $t$  and the number of bees increase. That is, a  $\frac{p}{p+q}$ -part of the bee population will be at  $Y$ , while the others, a  $\frac{q}{p+q}$ -part, will be at  $B$  for large  $t$  and a large number of bees. In other words, one would observe the ideal free distribution.

## IFD with a continuum of bees

Consider a continuum of identical bees that are all using the failures strategy  $A(r, s)$  with  $r, s \in \mathbb{N}$ , to forage on patches  $Y$  and  $B$  that have respectively total nectar supply  $y$  and  $b$ . The quantities are shared at each color

by the present bees. Given critical levels for the bees at  $Y$  and  $B$ , the “full-flower” probabilities,  $p$  and  $q$  respectively, are determined by the proportions of bees currently present in each of the patches. This population of bees can be distributed according to the IFD, i.e. there are full-flower probabilities  $p$  and  $q$ , and related critical levels, for  $Y$  and  $B$  respectively, such that each bee using  $A(r, s)$  is matching  $p/q$ , while the ratio of the fractions of bees at  $Y$  and  $B$  is  $y/b$ . More formally:

**Theorem 3.5** *If  $y > b$  and  $ys > br$ , then there exist  $p, q \in (0, 1)$  such that*

$$\frac{y}{b} = f(r, s) = \frac{q\bar{q}^s(1 - \bar{p}^r)}{p\bar{p}^r(1 - \bar{q}^s)} = \frac{p}{q}. \quad (12)$$

**Proof:** For  $q \in (0, b/y]$  define:

$$g(q) = \frac{(y/b)^2(1 - qy/b)^r(1 - (1 - q)^s)}{(1 - q)^s(1 - (1 - qy/b)^r)} \quad (13)$$

Then  $g$  is continuous on  $(0, b/y]$  with  $g(b/y) = 0$  and with

$$\begin{aligned} \lim_{q \downarrow 0} g(q) &= (y/b)^2 \lim_{q \downarrow 0} \frac{1 - (1 - q)^s}{1 - (1 - qy/b)^r} \\ &= (y/b)^2 \lim_{q \downarrow 0} \frac{s(1 - q)^{s-1}}{(ry/b)(1 - qy/b)^{r-1}} \\ &= \frac{(y/b)^2 s}{ry/b} = \frac{ys}{rb} > 1. \end{aligned} \quad (14)$$

By continuity of  $g$  there exists  $q^* \in (0, b/y]$  with  $g(q^*) = 1$ . Now  $q^*$  and  $p^* = yq^*/b$  are as desired.  $\square$

Note that this theorem is a static result for the existence of IFD with matching bees doing  $A(r, s)$ . The result does not provide a dynamic process of how to reach the IFD. We have to assume a continuum of bees in order to have  $p$  and  $q$  not being affected by single bees moving from one state of the automaton to another state of the automaton. The proportions of bees in the states of the automaton have to be independent of time. With a finite number of bees the probabilities  $p$  and  $q$  would always depend on the precise number of bees in those states.

## Attitude Towards Risk

Consider the single-bee situation of foraging on two patches  $Y$  and  $B$ , where all flowers have a normally distributed nectar supply with a common mean  $\mu$ . The yellow flowers have variance  $y$ , the blue ones have variance  $b$ .

**Theorem 3.6** *Suppose the bee is using a failures strategy and has a fixed critical level  $cl$ . Without loss of generality assume  $y < b$ .*

*If  $cl < \mu$ , then the bee will act as if it is risk averse, i.e. it will spend more time on  $Y$  than on  $B$ .*

*If  $cl > \mu$ , then the bee will act as if it is risk prone, i.e. it will spend more time on  $B$  than on  $Y$ .*

**Proof:** If  $cl < \mu$ , then the probability of getting a full flower is larger for  $Y$  than for  $B$  (Figure 2 in Section 2). Hence the probability  $p$  of getting a full  $Y$  flower is larger than  $q$  for a  $B$  flower. Since the bee's mechanism will find an automaton to match  $p$  and  $q$ , the bee will spend more time in  $Y$  than in  $B$ . The second part is similar.  $\square$

Remark that this result can also be extended to apply to other nectar distribution functions, like the ones discussed for the  $\varepsilon$ -sampling strategy.

Finally for this section we like to remark that we have only considered finite automata of the type: leave  $Y$  ( $B$ ) after  $y$  ( $b$ ) consecutive failures. However, one could also examine automata of a more general type. Let  $S, S_Y$  and  $S_B$  be finite non-empty sets of states with  $S = S_Y \cup S_B$  and  $S_Y \cap S_B = \emptyset$ . We can describe a foraging automaton for the bee as a map  $T : S \times \{0, 1\} \rightarrow S$  with the interpretation that a bee in state  $s \in S_Y$  is visiting a yellow flower; if this particular yellow flower is full (empty) then the bee moves to state  $T(s, 1)$  (resp.  $T(s, 0)$ ). Further research is required to fully understand the possibilities of obtaining matching and IFD results by means of these general type foraging automata.



## 4 Discussion and Related Literature

### The Matching Law

Many studies, articles, and even special books have been devoted to the phenomena of "the Matching Law." (Some main and recent references are Simon 1956, Herrnstein 1970, Heyman 1979, Commons et al. 1982, Houston et al. 1982, Houston 1983, Staddon 1983, Davison & McCarthy 1988, Staddon & Horner 1989). The reader of the above literature will probably be confused (as we are) by the rich and complicated details of the experimental results as well as by the theoretical analyses. There are very many versions of the matching law mathematical function (compare for example Krebs & Kacelnik 1991 to Houston & Sumida 1987 to Davison & McCarthy 1988 to the commentary by Herrnstein & Vaughan in Maynard Smith 1984).

In this study we focussed on one main issue related to the matching law: why the animal in an artificial binary choice setting behaves "irrationally" and does not go exclusively to the resource with the higher probability? Instead, the matching law reveals (Krebs & Kacelnik 1991) that the animal allocates its long-run behavior to two alternatives in proportion to the reward it obtains from them. To answer this question, we presented boundedly rational strategies for which the irrational matching behavior is compatible with the equilibrium conditions of the natural IFD.

We have presented foraging rules that lead the forager to match over time the reward probabilities by the fractions of the visits to the alternatives. However, one should not confuse this matching law with "probability matching" (Maynard Smith 1984). Probability matching is a foraging rule where, at each point in time, the forager is choosing an alternative according to the observed full flower probabilities at the respective patches. While probability matching is a moment-to-moment foraging rule, the matching

law describes the average behavior of the forager over an asymptotically long period of time.

## Learning Rules

As Maynard Smith (1984) points out, a learning rule (i.e. a foraging strategy) that we observe in animals should have several properties:

- a. It should have the *relative payoff sum* (RPS) property, which says that, after a sufficiently long period of time, the probability of doing a certain act should equal the total payoff received so far for doing this act, divided by the total payoff received so far for all acts.
- b. None of the available acts should ever fall to zero probability, since the environment might change.
- c. Any naive animal should start with some prior probabilities of performing the different acts.
- d. Recent payoffs should have a bigger effect on behavior than early ones (discount factor).

A strategy having these properties is Harley's RPS rule (Harley 1981, Maynard Smith 1984). For a two-choices situation  $(Y, B)$  this rule is defined as follows (Harley 1981):

Let  $r_Y, r_B > 0$  (residual values) and let  $0 < x < 1$  (memory factor). Let  $P_i(t)$  denote the payoff in  $i \in \{Y, B\}$  at time  $t \in \mathbb{N}$ . Now let  $S_i(0) = r_i$  and for  $t \geq 1$  define

$$S_i(t) = xS_i(t-1) + (1-x)r_i + P_i(t). \quad (15)$$

At time  $t$  choose alternative  $i \in \{Y, B\}$  with probability

$$f_i(t) = \frac{S_i(t-1)}{S_Y(t-1) + S_B(t-1)} \quad (16)$$

"In words, the (this) RPS rule says the following: display most frequently the behaviour which has, up to the present, paid the most, but only in proportion, roughly, to its cumulative payoff relative to the overall total" (Harley 1981). Although this rule is fairly simple, the foraging animal is required to adjust its probabilities of choosing either *Y* or *B* at each point in time. Compared to it we have presented an explicit "molecular" ("moment to moment" in Krebs & Kacelnik 1991 terms) strategy that has the above properties, but for which the foraging animal will only switch every once in a while. Notice that Harley's rule assumes the animal to update probabilities at all time points and chooses *Y* or *B* independent of the patch it is currently visiting. We do not believe that bees are capable of updating and computing probabilities before each visit (10 to 40 visits per minute!). Our foraging strategies give the same results while the bee is following very simple movement rules.

## Milinski Fishes

Our  $\epsilon$ -sampling strategy explains explicitly the dynamics in which "Milinski fishes" reach the IFD (Milinski 1979, 1984, Godin & Keenleside 1984, Milinski & Parker 1991). Each fish has to remember only an estimate of the average payoff received so far in the old resource, and to compare it with the current payoff in the new food source when sampling. If the current payoff is higher, the fish stays; if it is lower, it goes back to the old food source. This biological behavior corresponds to the potential function of Monderer & Shapley (1988). Milinski & Parker (1991, page 144) have a stationary model of the IFD with foraging fishes, but they do not have an explicit dynamic model by which the fishes can reach the IFD. Their model (including Parker & Sutherland 1986) can be viewed as a special case of the short-run stable matching of Peleg & Shmida (1992) where the fishes' different competitive weights correspond to the bees' different handling times

(technological abilities, sensu Selten 1978).

Godin & Keenleside (1984) have shown experimentally that the IFD can be achieved through "sampling." However, no explicit strategy has been suggested in the literature of how fish carried out the sampling procedure. Our  $\epsilon$ -sampling strategy corresponds to the "Major-Minor behavior" of bees, which has been observed in bumblebees by Heinrich (1979). The bumblebee visits mainly a certain flower type, the major one, but once in a while it samples (doing minoring) other flower types. It would be interesting to investigate how an individual bee's parameter  $\epsilon$  is influenced by its life history and its environment.

Milinski (1984) and Godin & Keenleside (1984) received interesting results when comparing the switching rate between two resources of fish with different competitive ability: Individuals experiencing a high feeding rate (high payoff) tend to switch patches less frequently than those individuals that received lower payoffs. These results can be explained by our  $\epsilon$ -sampling strategy (and also by the failure strategy). The fishes which are receiving less, reach the threshold of leaving more frequently, and the initiative to sample somewhere else is greater when  $\epsilon$ -sampling is payoff dependent (with  $\epsilon$  increasing with low payoffs and decreasing with high payoffs).

## Attitude Towards Risk

Our model explains in a simple way (without the need of utility theory) the well-known pattern in animal behavior of risk aversion and risk proneness (Real & Caraco 1986, Stephens & Krebs 1986, Krebs & Kacelnik 1991, McNamara & Houston 1992): In an environment in which the resource is normally distributed, the critical level can be interpreted as an existing condition for the animal. If the critical level is below the resource mean (Figure 2), then the animals should behave as if they are risk averse, while



if the critical level is above the mean, then they should behave as if they are risk prone. Our model predicts only a tendency to visit one of the resources more frequently and not to make an exclusive choice for one of the resources. In the case where nectar is distributed as in Figure 3, no matter what its critical level is, the bee will prefer the blue flower.

The elucidated review on risk sensitivity by McNamara & Houston (1992) has technically similar results to our points on attitude towards risk. However, they ask themselves which alternative the animal would prefer, while we find out what would happen to the animal if we take into account the observations of IFD and matching. In their model the critical level is a kind of evolutionary knowledge which, in a sense uses complete information about the environment. In our model the animal's actions are governed only by its own recent experiences and no complete information is required.

## Bounded Rationality and Myopic Learning

Our approach to studying foraging behavior is quite different from the "optimal foraging theory" used in ecology (Pyke et al. 1977, Krebs & Kacelnik 1991, Stephens & Krebs 1986, Bernstein et al. 1988). In optimal foraging theory the decision to leave a patch (resource alternative) is based on a comparison to the surroundings, in other words, it assumes that the animal has complete information and a powerful memory and computational ability. Our basic approach is that the animal uses "bounded memory" and makes its decisions only on the basis of its own recent experience. It does not know its competitor's moves/payoffs. It even may not know that it is involved in a game situation. The animal uses very simple decision rules to decide when to leave a patch (flower type), to decide where to go, and to decide whether or not to stay at the "new" patch. In such very fast biological activities (one bee makes very many visits per minute) the

animals use mainly their short term memory (Menzel 1985, 1990) and the movement rules depend on very short recall (e.g. remember the last 1-3 flowers) and very simple calculations (Real et al. 1990, Real 1991, 1992). These issues are related to myopic learning models (Monderer & Shapley 1988, Fudenberg & Kreps 1991, Milgrom & Roberts 1991, Monderer & Sela 1993) and to theory on bounded rationality (Kalai 1988, Selten 1990, Aumann 1992). We believe that bees and most other animals are not able to do sophisticated calculation of probabilities (e.g. as in the RPS-rule of Harley (1981), the learning rule by McNamara & Houston (1985) or in learning rules that require bayesian updating) and can neither update expected payoffs by functions that comprise long recall of history. Of course, in a matching experiment the reward probabilities are not known to the bee; these are known only to the researcher. One of the main limitations of many of the previous models is that they analyze the situation as if the bee does know those probabilities.

This bounded rationality approach is explained nicely in Boyd & Richerson 1985 (p. 93) when they address the issue of how animals and people apply Bayesian rationality: "simple rules of thumb (... called heuristics by psychologists) may greatly reduce cognitive complexity of decisions but still result in behavior that closely approximates normatively rational behavior in some restricted range of environments ... These heuristics often work well but occasionally lead to behavior that is irrational according to canons of Bayesian rationality."

In the foregoing pages we have presented an explicit model in which simple decision rules, that need only very short recall and only elementary calculations, can dictate an optimal foraging strategy. We have seen that finite automata with very few states appear to be appropriate tools for such tasks (Ben-Porath 1991, Ben-Porath & Peleg 1987). Our model stands in agreement with recent experiments of foraging animals which reveal that

animals take into account only the last 1-3 rewards when making short-term decisions (sensu Menzel 1990). It also fits the "melioration approach" (Herrenstein & Vaughan 1980) that animals use very simple choice rules which lead them to behave (sub)optimally in very complex natural situations (Houston et al. 1982, Hinson & Staddon 1983 and Boyd & Richerson 1985). To an outside observer this simple behavior looks very sophisticated and complicated as if they use utility functions, probability updating and complicated calculus. We believe that the same approach can explain some "irrational" behavior in economics and sociology; some "bounded rational" - ad hoc - strategies that are optimal under particular natural conditions may look very irrational in artificial situations. This is illustrated by the comparison of the matching experiment and the IFD phenomenon in nature.

## Acknowledgements

We would like to thank the Center for Rationality at the Hebrew University of Jerusalem for providing us with the environment to work together. Valuable comments by S. Ellner, J. Kagel and R. Selten are gratefully acknowledged. Moreover, A.S. would like to thank colleagues who helped him to enter the "orchard of mathematical economics": Dov Monderer for the potential games, Dov Samet and A. Neyman for finite automata, and Joel Sobel, A. Roth, and R. Selten for many hours of discussion on flowers, bees, rationality and game theory.

## Bibliography

- Aumann, R.J. (1992) Irrationality in Game Theory. In: *Economic analysis of Markets and Games, Essays in Honor of Frank Hahn*, P. Dasgupta, D. Gale, O. Hart, and E. Maskin (eds.). MIT Press, Cambridge, pp 214-227.
- Ben-Porath, E. (1991) Repeated games with finite automata. Working paper No. 7. The Suckler Institute of Economic Studies. Tel Aviv University.
- Ben-Porath, E. & B. Peleg (1987) On the folk theorem and finite automata. Center for Research in Mathematical Economics and Game Theory. Mimeo.
- Bernstein, C., A. Kacelnik & J.R. Krebs (1988) Individual decisions and the distribution of predators in a patchy environment. *J. Anim. Ecol.* 57:1007-1026.
- Boker, G. (1993) Pollination ecology of *Eremostachys laciniata* (Lamiaceae). M.Sc. Thesis, Hebrew University of Jerusalem, Dept. of Horticulture, Faculty of Agriculture.
- Boyd, R. & P.J. Richerson (1985) *Culture and the Evolutionary Process*. The University of Chicago Press, Chicago.
- Bush, R.R. & F. Mosteller (1955) *Stochastic Models for Learning*. Wiley, New York.
- Commons, M.L., R.J. Herrnstein & H. Rachlin (1982) *Quantitative Analysis of Behavior*, Vol. 2. *Matching and Maximizing Accounts*. Ballinger, Cambridge, Mass.
- Cresswell, J.E. (1990) How and why do nectar-foraging bumblebees initiate movements between inflorescences of wild bergamon *Monarda fistulosa* (Lamiaceae)? *Oecologia* 82: 450-460.
- Davison, M. & D. McCarthy (1988) *The Matching Law*. Lawrence Erlbaum Associates, Hillsdale, N.J.

- DeGroot, M. (1970) Optimal Statistical Decisions. McGraw-Hill, New York.
- Fishburn, P.C. (1976) Continua of stochastic dominance relations for bounded probability distributions. *J. of Math. Economics* 3:295-312.
- Fretwell, S.D. (1972) Seasonal Environments. Princeton Univ. Press, Princeton, New Jersey.
- Friedman, J. & A. Shmida (1992) Pollination, gathering nectar, and the distribution of flower species. Discussion paper no. 14, Center for Rationality, The Hebrew University of Jerusalem.
- Fudenberg, D. & D. Kreps (1991) Learning and equilibrium in games. Mimeo.
- Godin, J.G. & M.H.A. Keenleside (1984) Foraging on patchily distributed prey by a cichlid fish: a test of the Ideal Free Distribution theory. *Animal Behavior* 32:120-131.
- Hammerstein, P. & R. Selten (1993) Evolutionary game theory. In: *Handbook of Game Theory with Economic Applications*, R.J. Aumann & S. Hart (eds.). North-Holland.
- Harder, L.D. (1987) Choice of individual flowers by bumble bees: Interaction of morphology, time and energy. *Behavior* 104:60-77.
- Harley, C.B. (1981) Learning the evolutionary stable strategy. *J. Theor. Biol.* 89:611-633.
- Heinrich, B. (1979) Bumblebee Economics. Harvard Univ. Press, Cambridge, Mass.
- Herrenstein, R.J. (1970) On the law of effect. *J. Exp. Ana. Behavior* 13:243-266.
- Herrenstein, R.J. & W. Vaughan Jr. (1980) Melioration and behavioral allocation. In: *Limits to Action*, J.E.R. Staddon (ed.). Academic Press, New



York.

Heyman, G.M. (1979) Markov model description of changeover probabilities on concurrent variable interval schedules. *J. Exp. Ana. Behavior* 31:41-51.

Hinson, J.M. & J.E.R. Staddon (1983) Hill-climbing by pigeons. *J. Exp. Ana. Behavior* 39:25-47.

Houston, A.I. (1983) Optimality theory and matching. *Behavior Analysis Letters* 3:1-15.

Houston, A.I., A. Kacelnik, & J.M. McNamara (1982) Some learning rules for acquiring information. In: *Functional Ontogeny*, D.J. McFarland (ed.). Pitman, New York.

Houston, J.M. & B.H. Sumida (1987) Learning rules, matching and frequency dependence. *J. Theor. Biol.* 126:289-308.

Kadmon, R. & A. Shmida (1992) Departure rules used by bees foraging for nectar: A field test. *Evo. Eco.* 6:142-151.

Kadmon, R., A. Shmida, & R. Selten (1992) Within-plant foraging behavior of bees and its relationship to nectar distribution in *Anchusa strigosa*. *Isr. J. Bot.* 40:283-294.

Kalai, E. (1988) Bounded rationality and strategic complexity in repeated games. In: *Game Theory and Applications*, T. Ichiishi, A. Neyman & Y. Tauman (eds.). Academic Press, New York, pp 131-157.

Krebs, J.R., A. Kacelnik, & P. Taylor (1978) Test of optimal sampling by foraging great tits. *Nature* 275:27-31.

Krebs, J.R. & A. Kacelnik (1991) Decision-making. In: *Behavioral Ecology*, J.R. Krebs & N.B. Davies (eds.). Blackwell Scientific Pub., London, pp 105-136.

Laverty, T.M. (1980) The flower-visiting behavior of bumble bees: floral complexity and learning. *Can. J. Zool.* 58:1324-1335.

- Maynard Smith, J. (1982) *Evolution and the Theory of Games*. Cambridge University Press, Cambridge.
- Maynard Smith, J. (1984) Game theory and the evolution of behavior. *The Behavioral and Brain Sciences* 7:95-125.
- McNamara, J.M. & A.I. Houston (1985) Optimal foraging and learning. *J. Theor. Biol.* 117:231-249.
- McNamara, J.M. & A.I. Houston (1992) Risk-sensitive foraging: A review of the theory. *Bulletin of Mathematical Biology* 54:355-378.
- Menzel, R. (1985) Learning in honey bees in an ecological and behavioral context. In: *Experimental Behavioral Ecology*, Holldobler & Lindauer (eds.). G. Fischer Verlag, Stuttgart New York, pp 55-73.
- Menzel, R. (1990) Learning, memory and "cognition" in honey bees. In: *Neurobiology of Comparative Cognition*, R.P. Kesner & D.S. Olten (eds.). Erlbaum, Hillsdale, NJ, pp 237-292.
- Menzel, R. & U. Gregger (1992) Temporal dynamics and foraging behavior in honeybees. In: *Biology and Evolution of Social Insects*, J. Billen (ed.). Leuven University Press, Leuven, Belgium, pp 303-318.
- Menzel, R. & A. Shmida (1992) The ecology of flower colors and the natural color vision systems of insect pollinators. *Biol. Rev.* 68:81-120.
- Milgrom, P. & J. Roberts (1991) Adaptive and sophisticated learning in repeated normal form games. *Games and Eco. Beh.* 3:82-100.
- Milinski, M. (1979) An evolutionary stable feeding strategy in sticklebacks. *Z. Tierpsychol.* 51:36-40.
- Milinski, M. (1984) Competitive resource sharing: an experimental test of a learning rule for ESS's. *Animal Behavior* 32:233-242.
- Milinski, M. & G. Parker (1991) Competition for resources. In: *Behavioral Ecology*, J.R. Krebs & N.B. Davies (eds.). Blackwell Sc. Pub., Oxford, pp

137-168.

Monderer, D. & L.S. Shapley (1988) Potential Games. Mimeo.

Monderer, D. & A. Sela (1993) Fictitious play and no-cycling conditions. Mimeo.

Motro, U. & A. Shmida (1992) Near-far search: an evolutionary stable foraging strategy. Discussion paper no. 18, Center for Rationality, The Hebrew University of Jerusalem.

Neyman, A. (1985) Bounded complexity justifies cooperation in the finitely repeated prisoner's dilemma. *Economics Letters* 19:227-230.

Parker, G.A. & W.J. Sutherland (1986) Ideal free distributions when individuals differ in competitive ability: phenotype-limited ideal free models. *Anim. Behav.* 34:1222-1242.

Peleg, B. & A. Shmida (1992) Short-run stable matchings between bees and flowers. *Games and Econ. Beh.* 4:232-251.

Pyke, G.H., H.R. Pulliam, & E.L. Charnov (1977) Optimal foraging: A selective review of theory and tests. *The Quarterly Review of Biology* 52:137-154.

Real, L.A. (1991) Animal choice behavior and the evolution of cognitive architecture. *Science* 253:980-986.

Real, L.A. (1992) Information processing and evolutionary ecology of cognitive architecture. *Am. Nat.* 140:S108-S145.

Real, L. & T. Caraco (1986) Risk and foraging in stochastic environments. *Ann. Rev. Ecol. Syst.* 17:371-390.

Real, L.A., S. Ellner & L.D. Harder (1990) Short-term energy maximization and risk-aversion in bumble bees: A reply to Possingham et al. *Ecology* 71:1625-1628.



- Rothschild, M. (1974) A two-armed bandit theory of market pricing. *Journal of Economic Theory* 9:185-202.
- Selten, R. (1975) Re-examination of the perfectness concept for equilibrium points in extensive games. *International Journal of Game Theory* 4:25-55.
- Selten, R. (1978) A model of oligopolistic size structure profitability. *European Economic Review* 12:1-24.
- Selten, R. (1990) Bounded Rationality. *J. of Instit. and Theor. Eco.* 146:649-658.
- Shmida, A., C. O'Toole & Y. Ivri (1993) Elements of bee-flower pollination market in Israel. Mimeo.
- Shreiber, E. (1993) Foraging strategies in carpet inflorescences of *Alkanna strigosa* (Boraginaceae) by solitary bees. M.Sc. Thesis, The Hebrew University of Jerusalem. Dept. of Ecology and Evolution.
- Simon, H.A. (1956) A comparison of game theory and learning theory. *Psychometrika* 21:267-272.
- Staddon, J.E.R. (1983) *Adaptive Behavior and Learning*. Cambridge University Press, Cambridge.
- Staddon, J.E.R. & J.M. Horner (1989) Stochastic choice models: A comparison between Bush-Mosteller and a source-independent reward-following model. *J. Exp. Ana. Behavior* 52:57-64.
- Stephens, D.W. & J.R. Krebs (1986) *Foraging Theory*. Princeton University Press, Princeton, New Jersey.

**Discussion Paper Series, CentER, Tilburg University, The Netherlands:**

(For previous papers please consult previous discussion papers.)

<b>No.</b>	<b>Author(s)</b>	<b>Title</b>
9239	H. Bloemen	A Model of Labour Supply with Job Offer Restrictions
9240	F. Drost and Th. Nijman	Temporal Aggregation of GARCH Processes
9241	R. Gilles, P. Ruys and J. Shou	Coalition Formation in Large Network Economies
9242	P. Kort	The Effects of Marketable Pollution Permits on the Firm's Optimal Investment Policies
9243	A.L. Bovenberg and F. van der Ploeg	Environmental Policy, Public Finance and the Labour Market in a Second-Best World
9244	W.G. Gale and J.K. Scholz	IRAs and Household Saving
9245	A. Bera and P. Ng	Robust Tests for Heteroskedasticity and Autocorrelation Using Score Function
9246	R.T. Baillie, C.F. Chung and M.A. Tieslau	The Long Memory and Variability of Inflation: A Reappraisal of the Friedman Hypothesis
9247	M.A. Tieslau, P. Schmidt and R.T. Baillie	A Generalized Method of Moments Estimator for Long- Memory Processes
9248	K. Wärneryd	Partisanship as Information
9249	H. Huizinga	The Welfare Effects of Individual Retirement Accounts
9250	H.G. Bloemen	Job Search Theory, Labour Supply and Unemployment Duration
9251	S. Eijffinger and E. Schaling	Central Bank Independence: Searching for the Philosophers' Stone
9252	A.L. Bovenberg and R.A. de Mooij	Environmental Taxation and Labor-Market Distortions
9253	A. Lusardi	Permanent Income, Current Income and Consumption: Evidence from Panel Data
9254	R. Beetsma	Imperfect Credibility of the Band and Risk Premia in the European Monetary System
9301	N. Kahana and S. Nitzan	Credibility and Duration of Political Contests and the Extent of Rent Dissipation

<b>No.</b>	<b>Author(s)</b>	<b>Title</b>
9302	W. Güth and S. Nitzan	Are Moral Objections to Free Riding Evolutionarily Stable?
9303	D. Karotkin and S. Nitzan	Some Peculiarities of Group Decision Making in Teams
9304	A. Lusardi	Euler Equations in Micro Data: Merging Data from Two Samples
9305	W. Güth	A Simple Justification of Quantity Competition and the Cournot-Oligopoly Solution
9306	B. Peleg and S. Tijs	The Consistency Principle For Games in Strategic Form
9307	G. Imbens and A. Lancaster	Case Control Studies with Contaminated Controls
9308	T. Ellingsen and K. Wärneryd	Foreign Direct Investment and the Political Economy of Protection
9309	H. Bester	Price Commitment in Search Markets
9310	T. Callan and A. van Soest	Female Labour Supply in Farm Households: Farm and Off-Farm Participation
9311	M. Pradhan and A. van Soest	Formal and Informal Sector Employment in Urban Areas of Bolivia
9312	Th. Nijman and E. Sentana	Marginalization and Contemporaneous Aggregation in Multivariate GARCH Processes
9313	K. Wärneryd	Communication, Complexity, and Evolutionary Stability
9314	O.P. Attanasio and M. Browning	Consumption over the Life Cycle and over the Business Cycle
9315	F. C. Drost and B. J. M. Werker	A Note on Robinson's Test of Independence
9316	H. Hamers, P. Borm and S. Tijs	On Games Corresponding to Sequencing Situations with Ready Times
9317	W. Güth	On Ultimatum Bargaining Experiments - A Personal Review
9318	M.J.G. van Eijs	On the Determination of the Control Parameters of the Optimal Can-order Policy
9319	S. Hurkens	Multi-sided Pre-play Communication by Burning Money
9320	J.J.G. Lemmen and S.C.W. Eijffinger	The Quantity Approach to Financial Integration: The Feldstein-Horioka Criterion Revisited
9321	A.L. Bovenberg and S. Smulders	Environmental Quality and Pollution-saving Technological Change in a Two-sector Endogenous Growth Model

No.	Author(s)	Title
9322	K.-E. Wärneryd	The Will to Save Money: an Essay on Economic Psychology
9323	D. Talman, Y. Yamamoto and Z. Yang	The $(2^{n+m+1} - 2)$ -Ray Algorithm: A New Variable Dimension Simplicial Algorithm For Computing Economic Equilibria on $S^n \times R^m$
9324	H. Huizinga	The Financing and Taxation of U.S. Direct Investment Abroad
9325	S.C.W. Eijffinger and E. Schaling	Central Bank Independence: Theory and Evidence
9326	T.C. To	Infant Industry Protection with Learning-by-Doing
9327	J.P.J.F. Scheepens	Bankruptcy Litigation and Optimal Debt Contracts
9328	T.C. To	Tariffs, Rent Extraction and Manipulation of Competition
9329	F. de Jong, T. Nijman and A. Röell	A Comparison of the Cost of Trading French Shares on the Paris Bourse and on SEAQ International
9330	H. Huizinga	The Welfare Effects of Individual Retirement Accounts
9331	H. Huizinga	Time Preference and International Tax Competition
9332	V. Feltkamp, A. Koster, A. van den Nouweland, P. Borm and S. Tijs	Linear Production with Transport of Products, Resources and Technology
9333	B. Lauterbach and U. Ben-Zion	Panic Behavior and the Performance of Circuit Breakers: Empirical Evidence
9334	B. Melenberg and A. van Soest	Semi-parametric Estimation of the Sample Selection Model
9335	A.L. Bovenberg and F. van der Ploeg	Green Policies and Public Finance in a Small Open Economy
9336	E. Schaling	On the Economic Independence of the Central Bank and the Persistence of Inflation
9337	G.-J. Otten	Characterizations of a Game Theoretical Cost Allocation Method
9338	M. Gradstein	Provision of Public Goods With Incomplete Information: Decentralization vs. Central Planning
9339	W. Güth and H. Kliemt	Competition or Co-operation
9340	T.C. To	Export Subsidies and Oligopoly with Switching Costs
9341	A. Demirgüç-Kunt and H. Huizinga	Barriers to Portfolio Investments in Emerging Stock Markets

No.	Author(s)	Title
9342	G.J. Almekinders	Theories on the Scope for Foreign Exchange Market Intervention
9343	E.R. van Dam and W.H. Haemers	Eigenvalues and the Diameter of Graphs
9344	H. Carlsson and S. Dasgupta	Noise-Proof Equilibria in Signaling Games
9345	F. van der Ploeg and A.L. Bovenberg	Environmental Policy, Public Goods and the Marginal Cost of Public Funds
9346	J.P.C. Blanc and R.D. van der Mei	The Power-series Algorithm Applied to Polling Systems with a Dormant Server
9347	J.P.C. Blanc	Performance Analysis and Optimization with the Power-series Algorithm
9348	R.M.W.J. Beetsma and F. van der Ploeg	Intramarginal Interventions, Bands and the Pattern of EMS Exchange Rate Distributions
9349	A. Simonovits	Intercohort Heterogeneity and Optimal Social Insurance Systems
9350	R.C. Douven and J.C. Engwerda	Is There Room for Convergence in the E.C.?
9351	F. Vella and M. Verbeek	Estimating and Interpreting Models with Endogenous Treatment Effects: The Relationship Between Competing Estimators of the Union Impact on Wages
9352	C. Meghir and G. Weber	Intertemporal Non-separability or Borrowing Restrictions? A Disaggregate Analysis Using the US CEX Panel
9353	V. Feltkamp	Alternative Axiomatic Characterizations of the Shapley and Banzhaf Values
9354	R.J. de Groof and M.A. van Tuijl	Aspects of Goods Market Integration. A Two-Country-Two-Sector Analysis
9355	Z. Yang	A Simplicial Algorithm for Computing Robust Stationary Points of a Continuous Function on the Unit Simplex
9356	E. van Damme and S. Hurkens	Commitment Robust Equilibria and Endogenous Timing
9357	W. Güth and B. Peleg	On Ring Formation In Auctions
9358	V. Bhaskar	Neutral Stability In Asymmetric Evolutionary Games
9359	F. Vella and M. Verbeek	Estimating and Testing Simultaneous Equation Panel Data Models with Censored Endogenous Variables
9360	W.B. van den Hout and J.P.C. Blanc	The Power-Series Algorithm Extended to the <i>BMAP/PH/1</i> Queue



No.	Author(s)	Title
9361	R. Heuts and J. de Klein	An $(s,q)$ Inventory Model with Stochastic and Interrelated Lead Times
9362	K.-E. Wärneryd	A Closer Look at Economic Psychology
9363	P.J.-J. Herings	On the Connectedness of the Set of Constrained Equilibria
9364	P.J.-J. Herings	A Note on "Macroeconomic Policy in a Two-Party System as a Repeated Game"
9365	F. van der Ploeg and A. L. Bovenberg	Direct Crowding Out, Optimal Taxation and Pollution Abatement
9366	M. Pradhan	Sector Participation in Labour Supply Models: Preferences or Rationing?
9367	H.G. Bloemen and A. Kapteyn	The Estimation of Utility Consistent Labor Supply Models by Means of Simulated Scores
9368	M.R. Baye, D. Kovenock and C.G. de Vries	The Solution to the Tullock Rent-Seeking Game When $R > 2$ : Mixed-Strategy Equilibria and Mean Dissipation Rates
9369	T. van de Klundert and S. Smulders	The Welfare Consequences of Different Regimes of Oligopolistic Competition in a Growing Economy with Firm-Specific Knowledge
9370	G. van der Laan and D. Talman	Intersection Theorems on the Simplotope
9371	S. Muto	Alternating-Move Preplays and $vN - M$ Stable Sets in Two Person Strategic Form Games
9372	S. Muto	Voters' Power in Indirect Voting Systems with Political Parties: the Square Root Effect
9373	S. Smulders and R. Gradus	Pollution Abatement and Long-term Growth
9374	C. Fernandez, J. Osiewalski and M.F.J. Steel	Marginal Equivalence in $v$ -Spherical Models
9375	E. van Damme	Evolutionary Game Theory
9376	P.M. Kort	Pollution Control and the Dynamics of the Firm: the Effects of Market Based Instruments on Optimal Firm Investments
9377	A. L. Bovenberg and F. van der Ploeg	Optimal Taxation, Public Goods and Environmental Policy with Involuntary Unemployment
9378	F. Thuijsman, B. Peleg, M. Amitai & A. Shmida	Automata, Matching and Foraging Behavior of Bees

P.O. BOX 90153, 5000 LE TILBURG, THE NETHERLAND

**Bibliotheek K. U. Brabant**



17 000 01176359 7